



Short Communication

Monophyletic origin of brood care in damselfishes

Giacomo Bernardi*

Department of Ecology and Evolutionary Biology, University of California Santa Cruz, Santa Cruz, CA 95060, USA

ARTICLE INFO

Article history:

Received 13 October 2010

Revised 18 November 2010

Accepted 23 December 2010

Available online 25 January 2011

Keywords:

*Altrichthys**Acanthochromis*

Pomacentridae

Mitochondrial DNA

Nuclear DNA

Brood care

Formalin artifacts

ABSTRACT

The absence of a pelagic larval stage and brood care has evolved very few times in coral reef fishes. Damselfishes, a widely represented group with more than 380 species, includes only three such species, the monotypic *Acanthochromis polyacanthus*, and the two *Altrichthys* species, *Altrichthys azurelineatus* and *Altrichthys curatus*. In a recent study, Cooper et al. provided a comprehensive phylogenetic hypothesis for the damselfish family, based on mitochondrial and nuclear sequences, with more than 100 species that included all extant genera (Cooper et al., 2009). *A. polyacanthus* and sequences from formalin-preserved tissue of *A. curatus* did not cluster together, indicating that brood care may have evolved independently twice in damselfishes. Here, we use mitochondrial and nuclear molecular markers from fresh specimens of both *Altrichthys* species. We found that *Altrichthys* and *Acanthochromis* are closest relatives, which suggests that brood care may have evolved only once in damselfishes. Due to the limited geographic range of *Altrichthys* (western Philippines) and the abundant presence of *Acanthochromis* in that region, we also suggest that brood care may have originated in that region.

© 2011 Elsevier Inc. All rights reserved.

1. Introduction

Most reef fishes have a bipartite lifestyle that includes a sedentary adult stage and a pelagic larval stage that typically lasts from a few days to several weeks. Indeed, the lack of a pelagic larval stage is supposed to have a number of selective disadvantages including a difficulty in dispersing over great distances and its associated risk of local extinctions (Poethke et al., 2003), and very few species of marine fishes lack a pelagic larval stage (Doherty et al., 1985; Leis, 1991).

For species that lack a pelagic larval stage (apelagic species), gene flow was found to be limited over large geographic scales in the damselfish *Acanthochromis polyacanthus* (Doherty et al., 1994; Miller-Sims et al., 2008; Bay et al., 2008), the banggai cardinalfish, *Pterapogon kauderni* (Bernardi and Vagelli, 2004; Hoffman et al., 2005), and the surfperches *Embiotoca jacksoni* and *Embiotoca lateralis* (Bernardi, 2000, 2005). *A. polyacanthus* (Planes and Doherty, 1997; van Herwerden and Doherty, 2006) and *P. kauderni* (Vagelli et al., 2009) have also been shown to experience limited gene flow over smaller geographic scales. Strong founder effects, which are expected in apelagic species, have also been described in Coral Sea populations of *A. polyacanthus* (Planes et al., 2001). Therefore, while apelagic species are rare, they represent a unique opportunity to study adult dispersal, colonization dynamics, founder ef-

fects, speciation mechanisms, and population genetics issues in general.

The care of offspring, from newly hatched larvae all the way to the recruitment stage, referred here as brood care (*sensu*-Cooper et al., 2009) provides some distinctive evolutionary advantages. Besides direct benefits, such as the provision of food when young can feed on the parents' body mucus (Robertson, 1973; Allen, 1999), and the protection of the young from predation, indirect benefits include an increased capacity for local adaptation, and the advantages that the young obtain from being directly recruited into appropriate habitats by their parents. In coral reef fishes, however, apelagic strategies have evolved only rarely (Leis, 1991).

For example, all cardinalfish (Apogonidae) males keep eggs in their mouths (mouthbrooders), yet only two species, *Pterapogon kauderni* and *P. mirifica*, prolong the care by keeping the young in their mouths until ready for recruitment (Vagelli, 1999; Vagelli pers. com.). Damselfishes (Pomacentridae) include approximately 380 species that build nests and spawn benthically (Nelson, 2006). Eggs are fanned and protected by the parents, a preadaptation that seems conducive to further brood care. Yet only three species lack a pelagic larval stage: the spiny damselfish, *A. polyacanthus*, a relatively large species that is widely distributed from Indonesia and the Philippines to the Great Barrier Reef and the Solomon Islands, and two congeneric species, *Altrichthys azurelineatus* and *Altrichthys curatus*, that live mostly sympatrically in a very restricted area of western Philippines in the Calamian islands, and for *A. curatus*, in the Cuyo Islands. Individuals of the two *Altrichthys* species live in close proximity of each other, but tend to select dif-

* Address: Department of Ecology and Evolutionary Biology, University of California Santa Cruz, 100 Shaffer Road, Santa Cruz, CA 95060, USA.

E-mail address: bernardi@biology.ucsc.edu

ferent microhabitats, *A. curatus* prefers to associate with *Porites cylindrica* corals (Allen, 1999), while *A. azurelineatus* has less specific microhabitat preferences and is more generally distributed (Allen, 1999). In the Calamian islands, *A. azurelinatus* and *A. curatus* also live in very close proximity to *A. polyacanthus* and another damselfish, *Pomacentrus stigma* (pers. obs.).

A recent comprehensive phylogeny of the damselfish family that included all genera and more than 100 species, based on mitochondrial and nuclear markers, uncovered significant evolutionary characteristics for this group (Cooper et al., 2009). In this study, the genus *Altrichthys* was represented by formalin-preserved tissue of *A. curatus* collected in 1978 in the Cuyo Islands (USNM 00348456). Extracting DNA from formalin-preserved tissue is a technical challenge, but it is sometimes required when no other options are available. Based on mitochondrial and nuclear DNA sequences, data suggested that brood care evolved twice in damselfishes. Brood care evolved once with *A. polyacanthus*, which in this study was the sister taxon to *Hemiglyphidodon plagiometopon* (a species that does not show brood care, Hoey, pers. com.), and this clade was the sister lineage to the branch that contains the genera *Amblyglyphidodon* and *Neoglyphidodon*. Brood care evolved independently a second time with *A. curatus*, which grouped with *Pomacentrus stigma* and *Pomacentrus lepidogenys*. Thus, while the lack of a pelagic larval stage is extremely rare in coral reef fishes, it was hypothesized that such traits evolved independently at least twice in damselfishes (Cooper et al., 2009).

The goal of our study was to reassess this statement using fresh material for both *Altrichthys* species, in order to fully understand the geographic and evolutionary components of evolution of brood care in damselfishes.

2. Materials and methods

2.1. Collections and DNA samples

Four individuals of each *A. azurelineatus* and *A. curatus* were collected by hand nets in the islands of Coron (COR), Dibatuc (DIB), and Tangat (TAN), in the Calamian Archipelago, Palawan, Philippines. After collection, samples were immediately placed in 95% ethanol and stored at ambient temperature in the field, and then at 4 °C in the lab. Muscle tissue was later dissected from these samples. Total genomic DNA was prepared from 75 to 150 mg of muscle or liver tissue by proteinase K digestion in lysis buffer (10 mM Tris, 400 mM NaCl, 2 mM EDTA, 1% SDS) overnight at 55 °C. This was followed by purification using chloroform extractions and alcohol precipitation (Sambrook et al., 1989).

2.2. PCR amplification and sequencing

In order to compare our data with previously published sequences, we PCR amplified one mitochondrial marker, 16S rRNA, and two nuclear markers, RAG1 and RAG2. Primers and protocols followed Cooper et al. (2009). After purification of the PCR products, following the manufacturer's protocol (ABI, Perkin-Elmer), sequencing was performed with the primers used in the PCR amplification on an ABI 3100 automated sequencer (Applied Biosystems, Foster City, CA). In the case of the nuclear loci, all individuals were homozygous, which made the calling of alleles possible.

Additional sequences from Cooper et al. (2009) were used in our analyses. Sequences from formalin preserved *A. curatus* were limited to the 16SrRNA marker and for 232 bp of the RAG2 marker. All other sequences used here were for all three markers, 16SrRNA, RAG1, and the complete RAG2. The phylogeny presented by Cooper et al. (2009) comprises 104 species, which are mostly outside the subfamily (Pomacentrinae) that is germane to our study. We there-

fore decided to choose a subsample of the original dataset that is more pertinent to our question by selecting random representatives of all the major subclades that were found in topological proximity to *A. polyacanthus* and *A. curatus*. These are: *Premnas biaculeatus*, *A. polyacanthus*, *H. plagiometopon*, *Amblyglyphidodon leucogaster*, *Neoglyphidodon nigroris*, *Amblypomacentrus clarus*, *Pomacentrus amboinensis*, *Pomacentrus coelestis*, *P. lepidogenys*, *P. stigma*, and *Pomacentrus brachialis*. In the following phylogenetic analyses, the spinecheek clownfish, *P. biaculeatus*, was used as an outgroup, following Cooper et al. (2009).

2.3. Phylogenetic analyses

We used the computer program MAFFT (Katoh et al., 2002) implemented by the Geneious software package (Drummond et al., 2009) to align the DNA sequences. Phylogenetic relationships were assessed by Maximum Likelihood (ML, GARLI software, Zwickl, 2006), Maximum Parsimony (MP, PAUP* software Swofford, 2003), and Neighbor-Joining (NJ, PAUP software), methods. For Maximum Likelihood topologies, we conducted 10 independent runs in GARLI, using default settings and the automated stopping criterion, terminating the search when the ln score remained constant for 20,000 consecutive generations. The best likelihood of those runs was retained and is presented here (Fig. 1). Maximum Parsimony searches included 100 random addition replicates and TBR branch swapping with the Multrees option. Neighbor-Joining reconstructions used distances based on substitution models obtained with Modeltest (HKY + G). Statistical confidence in nodes was evaluated using 2000 non-parametric bootstrap replicates (Felsenstein, 1985) (100 replicates for Maximum Likelihood in GARLI, using the automated stopping criterion set at 10,000 generations). Topological differences were tested using a Shimodaira and Hasegawa test (Shimodaira and Hasegawa, 1999) implemented by PAUP, based on resampling of estimated log-likelihoods tests (RELL, 1000 replicates).

3. Results

3.1. Sequences

Mitochondrial 16SrRNA (484 bp) sequences, and nuclear RAG1 (900 bp) and RAG2 (766 bp) sequences were obtained from all of the individuals sampled. Out of an aligned total of 2152 bp, 312 bp were variable and 164 bp were phylogenetically informative. Sequences were deposited in GenBank (Accession Numbers HQ629877–HQ629900).

3.2. Phylogenetic relationships

Due to the technical difficulty of sequencing formalin preserved specimens, the published sequences of *A. curatus* are shorter than the sequences of other damselfishes. We therefore reconstructed phylogenetic relationships based on two datasets. The first dataset included the sequences of *A. curatus* from Cooper et al. (2009), and was therefore a truncated set of sequences. This included the full length of the 16SrRNA marker (484 bp) and a portion of the RAG2 marker (232 bp). The second dataset included the full length of sequences for the three markers, but did not include the *A. curatus* sample from Cooper et al. (2009).

Besides the presence or absence of the 2009 *A. curatus*, all three phylogenetic reconstruction methods based on either datasets resulted in topologies that were not statistically different. A phylogenetic tree based on a full dataset, but with an indicated position of the 2009 *A. curatus* individual, is presented in Fig. 1.

further work is warranted to elucidate the mechanisms that led to their divergence.

The third apelagic damselfish, *A. polyacanthus*, is markedly different from its closest relatives, *A. azurelineatus* and *A. curatus*. *A. polyacanthus* is a larger species that lives higher in the water column, two attributes that may have played a role in its much wider range and dispersal capability. Yet, since all three apelagic damselfishes are found sympatrically, it is tempting to think that the western Philippines may have been the center of origin for brood care in damselfishes.

Acknowledgments

I would like to thank James O'Donnell for help in the field, and James O'Donnell, Eric Crandall and Adam Hoey for helpful discussions.

References

- Allen, G.R., 1999. *Altrichthys*, a new genus of damselfish (Pomacentridae) from Philippine seas with description of a new species. *Rev. Fr. Aquariol.* 26, 23–28.
- Bay, L.K., Caley, M.J., Crozier, R.H., 2008. Meta-population structure in a coral reef fish demonstrated by genetic data on patterns of migration, extinction and recolonisation. *BMC Evol. Biol.* 8, 248.
- Bernardi, G., 2000. Barriers to gene flow in *Embiotoca jacksoni*, a marine fish lacking a pelagic larval stage. *Evolution* 54, 226–237.
- Bernardi, G., 2005. Phylogeography and demography of sympatric sister species, *Embiotoca jacksoni* and *E. lateralis* along the California coast: historical versus ecological factors. *Evolution* 59, 386–394.
- Bernardi, G., Vagelli, A., 2004. Population structure in Banggai cardinalfish, *Pterapogon kauderni*, a coral reef species lacking a pelagic larval phase. *Mar. Biol.* 145, 803–810.
- Cooper, W.J., Smith, L.L., Westneat, M.W., 2009. Exploring the radiation of a diverse reef fish family: phylogenetics of the damselfishes (Pomacentridae), with new classifications based on molecular analyses of all genera. *Mol. Phylogenet. Evol.* 52, 1–16.
- Doherty, P.J., Williams, D., Sale, P.F., 1985. The adaptive significance of larval dispersal in coral reef fishes. *Environ. Biol. Fish.* 12, 81–90.
- Doherty, P.J., Planes, S., Mather, P., 1994. *Acanthochromis polyacanthus*, a fish lacking larval dispersal, has genetically differentiated populations at local and regional scales on the Great Barrier Reef. *Mar. Biol.* 121, 11–21.
- Drummond, A.J., Ashton, B., Cheung, M., Heled, J., Kearse, M., Moir, R., Stones-Havas, S., Thierer, T., Wilson, A., 2009. Geneious v4.7. <<http://www.geneious.com/>>.
- Felsenstein, J., 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39, 783–791.
- Hoffman, E.A., Kolm, N., Berglund, A., Arguello, J.R., Jones, A.G., 2005. Genetic structure in the coral-reef-associated Banggai cardinalfish, *Pterapogon kauderni*. *Mol. Ecol.* 14, 1367–1375.
- Katoh, K., Misawa, K., Kuma, K., Miyata, T., 2002. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res.* 30, 3059–3066.
- Leis, J.M., 1991. The pelagic stage of Reef Fishes. In: Sale, P. (Ed.), *The Ecology of Fishes on Coral Reefs*. Academic Press, Inc., San Diego, CA, pp. 82–229.
- Miller-Sims, V., Gerlach, G., Atema, J., Kingsford, M.J., 2008. Dispersal in the spiny damselfish, *Acanthochromis polyacanthus*, a coral reef fish species without a larval pelagic stage. *Mol. Ecol.* 17, 5036–5048.
- Nelson, J.S., 2006. *Fishes of the World*, fourth ed. John Wiley & Sons, Hoboken, NJ.
- Planes, S., Doherty, P.J., 1997. Genetic and color interactions at a contact zone of *Acanthochromis polyacanthus*: a marine fish lacking pelagic larvae. *Evolution* 51, 1232–1243.
- Planes, S., Doherty, P.J., Bernardi, G., 2001. Strong genetic divergence among populations of a marine fish with limited dispersal, *Acanthochromis polyacanthus*, within the Great Barrier Reef and the Coral Sea. *Evolution* 55, 2263–2273.
- Poethke, H.J., Hovestadt, T., Mitesser, O., 2003. Local extinction and the evolution of dispersal rates: causes and correlations. *Am. Nat.* 161, 631–640.
- Robertson, D.R., 1973. Field observations on the reproductive behaviour of a pomacentrid fish, *Acanthochromis polyacanthus*. *Z. Tierpsychol.* 32, 319–324.
- Sambrook, J., Fritsch, E.F., Maniatis, T., 1989. *Molecular Cloning: A Laboratory Manual*, second ed. vol. 3. Cold Spring Harbor Laboratory Press, New York, 253p.
- Shimodaira, H., Hasegawa, M., 1999. Multiple comparisons of log-likelihoods with applications to phylogenetic inference. *Mol. Biol. Evol.* 16, 1114–1116.
- Swofford, D.L., 2003. PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods). Version 4. Sinauer Associates, Sunderland, Massachusetts.
- Vagelli, A.A., 1999. The reproductive biology and early ontogeny of the mouthbrooding Banggai cardinalfish, *Pterapogon kauderni* (Perciformes, Apogonidae). *Environ. Biol. Fish.* 56, 79–92.
- Vagelli, A., Burford, M.O., Bernardi, G., 2009. Fine scale dispersal in Banggai Cardinalfish, *Pterapogon kauderni*, a coral reef species lacking a pelagic larval phase. *Mar. Genom.* 1, 129–134.
- van Herwerden, L., Doherty, P.J., 2006. Contrasting genetic structures across two hybrid zones of a tropical reef fish, *Acanthochromis polyacanthus* (Bleeker 1855). *J. Evol. Biol.* 19, 239–252.
- Zhang, J., 2010. Exploiting formalin-preserved fish specimens for resources of DNA barcoding. *Mol. Ecol. Res.*, 935–941.
- Zwickl, D.J., 2006. Genetic Algorithm Approaches for the Phylogenetic Analysis of Large Biological Sequence Datasets under the Maximum Likelihood Criterion. Ph.D. Dissertation, The University of Texas at Austin.